West Chicago Urban Forest Summary

The Chicago Region Trees Initiative (CRTI) goal is that, by 2050, the Chicago Region will support and host a healthier urban forest, comprised of a diversity of tree species and ages, appropriately distributed across land use types in the region. The forest will provide the region improved environmental, economic, and social benefits. In order to achieve that goal CRTI works with a wide variety of people who work with and manage trees. This document is intended to help municipalities understand their urban forest, and identify strategies that they can use to make it better.

The *urban forest* is comprised of all of the trees in an urban setting, regardless of who owns or manages them. It is made up of street trees, forested natural areas and even the trees in resident's back yards. These trees are all included in the urban forest, because they all provide benefits that municipalities depend on. They improve air and water quality, reduce flooding and the urban heat island effect, and reduce energy use by shading buildings. Trees provide habitat for wildlife and improve residents' quality of life by reducing crime rates, increasing property value and boosting social cohesion in neigborhoods.

The magnitude of benefits that trees provide correlates with the size, structure and location of their canopy. Understanding the extent of tree canopy is critical for urban planning. Canopy maps can be used to quantify the benefits that their trees provide, identify where new plantings would have the greatest impact and to develop priorities and strategies for expanding the canopy.

The Chicago Region Trees Initiative, USDA Forest Service, American Forests, and the University of Vermont mapped land cover across the seven-county Chicago Region. This project not only identifies tree canopy, but also other green infrastructure including vegetation under 10 feet tall, bare soil and water; and gray infrastructure including buildings, roads and rail and other paved surfaces like sidewalks and parking lots (Fig. 1). Here after, these seven layers will be referred to as *land cover types*.



Fig. 1: Comparison of satellite image and land cover map. Seven types of gray and green infrastructure are in the land cover map.

West Chicago Urban Forest Summary



Fig 2: DuPage County's current land cover (A), including 26% canopy cover. An additional 54% of the county is suitable for planting (B). West Chicago currently has 15% canopy cover (C), and 68% of the land cover could potentially be converted to canopy (D).

Overall, 26% of DuPage County is covered by tree canopy (Fig. 2). There is a lot of room for growth across the county. We can identify spaces where trees could potentially be planted by adding together the vegetation, bare soil and other paved surface land cover types, as these land cover types could be converted to canopy with minimal effort. In all, these land cover types make up 54% of the county's area, meaning that canopy cover could potentially be raised to 83% if all of these surface were converted to trees. It is important to note, that while these surfaces could theoretically be covered with canopy, it is not necessarily preferable. Agricultural fields and baseball diamonds are included as "plantable space," but few would agree that these are ideal sites to expand the forest canopy.

These land cover data can also describe canopy at the municipal scale. West Chicago currently has 15% canopy cover, and could potentially increase their canopy to 83% (Fig. 2).

West Chicago Urban Forest Summary



Fig 3: Variations in land cover across land use types.

Canopy cover is not distributed evenly across the region, nor within municipalities. To better understand how land cover patterns vary, we can compare them across land use types, like residential, commercial or park properties. In West Chicago, the highest percentage of canopy is found in natural areas, cemeteries and residential properties (Fig. 3). Agriculture and water land use types have the lowest canopy cover. As one might expect, transit areas have a large proportion of roads, and residential and commercial land use types have an abundance of buildings. See Table 1 at the end of this report for more details.

West Chicago Urban Forest Summary



Fig 4: Current canopy and possible planting space across land use types.

By combining vegetation, bare soil and other paved surface categories we can identify which land use types have the most room for growth. In West Chicago, the highest proportions of plantable space are found in agriculture and utility sites (Fig. 4).



Fig 5: The majority of land is transit land use, followed by residential.

While utility properties have a high proportion of plantable space, these land use types make up a relatively small area in West Chicago (Fig. 5). The majority of its land is transit followed by residential.

West Chicago Urban Forest Summary



Fig 6: Transit has the greatest potential for increasing the canopy, followed by agriculture an residential.

Residential, transit, agricultural and commercial land use types have the most area that could possibly be converted to canopy (Fig. 6). Targeting these areas could have the greatest impact in expanding the canopy. However, each of these land use types will require different strategies to increase canopy. Many of the transit properties are publicly owned, and could therefore be the easiest to work with. Residential and commercial property owners could be encouraged to plant more trees through tree giveaways, ordinances that encourage tree preservation, or stormwater tax breaks for properties that have more tree canopy. While planting trees in agricultural properties may not always be feasible, there are ways to expand canopy in this land use type. Trees can be planted along road ways or drainage areas. Additionally, ordinances can mandate the planting of trees as agricultural sites are converted to other land use types.

Table 1: Summary of land cover across land use types.

	Tree canopy		Vegetation		Bare soil		Water		Buildings		Roads and rail		Other paved	
	Acres	*Percent*	*Acres*	*Percent*	*Acres*	*Percent*	*Acres*	*Percent*	*Acres*	*Percent*	*Acres*	*Percent*	*Acres*	*Percent*
Agriculture	70.7	5.7%	1111.0	89.9%	9.5	0.8%	22.2	1.8%	0.9	0.1%	1.1	0.1%	19.8	1.6%
Cemetery	6.0	34.1%	11.4	64.0%	0.0	0.0%	0.0	0.0%	0.0	0.0%	0.0	0.0%	0.3	1.9%
Commercial	119.1	8.9%	336.1	25.1%	16.8	1.3%	9.1	0.7%	349.5	26.1%	11.1	0.8%	495.9	37.1%
Golf	76.0	12.1%	490.2	78.0%	6.1	1.0%	32.4	5.2%	2.4	0.4%	6.9	1.1%	14.7	2.3%
Institutional	33.3	14.2%	119.2	50.7%	1.5	0.6%	0.8	0.3%	29.4	12.5%	1.1	0.5%	49.7	21.2%
Natural area	182.0	29.3%	414.5	66.7%	0.0	0.0%	22.0	3.5%	0.0	0.0%	0.4	0.1%	2.9	0.5%
Park	61.1	15.3%	273.4	68.5%	9.8	2.4%	38.2	9.6%	1.3	0.3%	1.7	0.4%	13.7	3.4%
Residential	484.5	29.1%	760.2	45.6%	2.0	0.1%	32.8	2.0%	254.1	15.3%	7.9	0.5%	124.2	7.5%
Transit	197.1	8.0%	1281.7	51.9%	12.4	0.5%	11.5	0.5%	28.4	1.2%	673.0	27.2%	267.2	10.8%
Utility	20.3	13.0%	113.4	72.9%	0.0	0.0%	3.6	2.3%	3.8	2.5%	0.4	0.3%	14.1	9.1%
Vacant	181.1	24.3%	510.2	68.4%	5.4	0.7%	22.3	3.0%	1.4	0.2%	4.3	0.6%	21.3	2.9%
Water	8.9	5.2%	79.3	46.0%	0.4	0.2%	73.8	42.8%	0.1	0.0%	1.9	1.1%	8.0	4.7%
Total abundance	1250.0	14.9%	4911.2	56.8%	58.2	0.7%	172.7	2.8%	669.8	6.9%	703.6	7.3%	1002.8	10.7%

West Chicago Urban Forest Summary

West Chicago has lower canopy than most of its neighbors (Fig. 7). Only Carol Stream has less.



Figure 7: Comparison of land cover of West Chicago and its neighbors.

West Chicago Urban Forest Summary



Figure 9: This Forest Service publication be found in its entirety at: http://www.nrs.fs.fed.us/pubs/44566.

While understanding canopy is an important component of urban forestry, canopy alone does not show the whole picture. We can use canopy to quantify the number and extent of trees, but not their identity, health, nor vulnerability to pests, diseases or climate change. In order to better understand the urban forest, we need a tree inventory, which will describe the abundance and location of tree species (Fig. 8).

Tree inventories come in many shapes and sizes. The most complete inventories gather data on every tree in the study area, and include information like each tree's species, any health issues the tree may have, and its specific location. This sort of inventory is invaluable for planning and monitoring the urban forest's health and growth over time. The Morton Arboretum and the USDA Forest Service conducted a sample inventory across the seven county region, and determined species composition at the county scale (Fig. 9). This resource is invaluable in understanding broad-scale composition.

To measure species composition on a finer scale, rely on inventories that measure every tree. Many communities, including West Chicago, have conducted such inventories on public property. We can use these inventories to better understand the state of its urban forest, and to compare municipalities who have an inventories to each other and to the region as a whole.

All calculations in this report were made using a 2015 inventory of West Chicago's municipal trees. It should be noted that this only represents a small portion of the trees in West Chicago, and a more compete inventory is necessary to further explore the urban forest.

Figure 8: Measuring tree size is a critical component to completing a tree inventory.

West Chicago Urban Forest Summary



Figure 10: Region-wide, maple is the most abundant genus by number of trees in municipal plantings (A), as well as by proportion of canopy (B). West Chicago has far more ashes than the rest of the region and fewer maples (C and D).

Inventories allow us to look at diversity in two ways, by the abundance of individual trees, and the proportion of the entire canopy that these trees make up. To determine the second figure, we can use the diameter of the trees as a correlate for canopy abundance, as those values are closely related. Both of these measures are important. Number of individual stems is useful when calculating the number of trees that might be affected by a given pest or disease. The abundance of canopy will show how the entire forest might change. That is, losing ten, small apple trees would have a much smaller impact on the ecosystem services that a forest offers than losing ten, mature oaks. Figure 10 shows the most abundant genera in West Chicago and across 55 other municipalities.

Many municipalities have very low species diversity. They rely heavily on tried and true species, like Freeman maples, white ash and honeylocust. Maples, on average, comprise 30% of municipal plantings. Ashes were generally in similar abundances, and losing them was a terrible ordeal. As devastating as losing ashes was for the Chicago region, losing maples would be much worse.

This illustrates how critical it is to actively increase species diversity where possible. Most pests and diseases (like emerald ash borer and Dutch elm disease) only attack a specific species or genera of plants. By diversifying species, we can ensure that our forest is resilient to these attacks. CRTI recommends that municipalities strive to have no more than 15% of a single family, 10% of a genus, and 5% of a species in their plantings. These guidelines should be met not only across the whole municipality, but at smaller scales like individual blocks.

West Chicago's urban forest is dominated by ashes. The town is likely in a state of transition as these trees are lost. While the town is replacing them, it is imperitive to increase species diversity so that the forest is resilient.



Figure 11: There are more trees in smaller age classes than in larger ones across all municipal inventories. West Chicago has fewer trees in the smallest size class, but otherwise follows this pattern as well.

There is another type of diversity to consider beyond species diversity. A sustainable forest has a variety of ages and sizes of trees. If all of the trees in an area were planted at the same time, they will grow, age and die at the same time. When these trees reach the end of their lives, it could leave a property without trees.

For that reason, it is important to try to increase age diversity of a forest. This can be done by planting trees over several years, planting trees with different growth habits (some trees grow quickly and have shorter lifespans than others), and by under planting aging trees, so that something is ready to replace them when they die.

Paying attention to size diversity is especially critical when recovering from emerald ash borer. Many communities have vowed to replace all of the trees that they remove, but this could become problematic if they are planting all of those trees within a short time period. It may be better for overall health of the forest to space these plantings out over several years.

Overall, size diversity averaged across all municipalities looks sustainable (Fig. 11). There is room for some trees to die between each age class, with plenty to remaining to grow into the next size class. West Chicago has fewer in the smallest size class. If this trend continues, canopy may decrease over time. This is especially worrisome, as much of West Chicago's canopy is made up of ashes, which are likely dying. It is imperitive to plant new trees to replace these dying ashes.

While size diversity is sustainable on the municipal, county and regional scale, it is also important to zoom into smaller areas, like specific subdivisions or properties. Planning for size class diversity on this level is important to local ecosystem services, like reducing energy use, managing stormwater and retaining soil.

West Chicago Urban Forest Summary

Urban trees are extremely valuable. Research has allowed us to to quantify the values that trees provide, and these values go far beyond the aesthetics that are readily recognized. For example:

- Urban trees save energy by reducing surface temperatures and shading buildings.
- They store carbon dioxide and remove pollutants from the air.
- They intercept stormwater and help reduce flooding.
- Residents preferentially buy properties that have more trees, meaning that trees increase property values.

The i-Tree suite of tools was developed by the US Forest Service. They allow users to calculate tree benefits at a variety of scales, from an individual tree, to entire tree inventories, to landscape scale assessments of canopy and hydrology. For more information on i-Tree tools and methodology visit iTreetools.org.

Figure 12 shows the benefits that all of the trees (including trees public and private property) in West Chicago offer. These values were calculated with i-Tree Landscape. Each year, West Chicago's trees provide the municipality with $884,300 worth of benefits. These trees also store a lot of carbon, which is valued at an additional $4,874,100.



Figure 12: Trees offer myriad benefits, including intercepting stormwater, improving air quality and removing carbon from the atmosphere.

West Chicago Urban Forest Summary

Cities tend to be hotter than rural areas because buildings and pavement absorb the sun's energy and release it as heat. This is known as the urban heat island effect. High urban temperatures increase the use of energy within buildings. It can also cause a variety of health issues to residents, and extreme heat can even cause death. Trees help lower urban temperatures by shading built surfaces and through evaporative cooling. Urban areas that have more tree canopy tend to have lower surface temperatures (Figure 13). Planting more trees in parking lots and around buildings can be especially helpful in reducing urban temperatures and making cities more comfortable.



Figure 13: The image on the left shows tree canopy and on the right shows surface temperature. Surface temperature was calculated using a landsat image from September 2014. Areas that have higher tree canopy tend to have lower temperatures.

West Chicago Urban Forest Summary



Figure 14: Oak ecosystems were abundant prior to Euro-American settlement. This map shows which of these ecosystems still remained in 1939 and 2010.

Oaks are a keystone species in our region's ecology. They provide habitat and food for countless animals, and they influence which plants grow around them. Prior to Euro-American settlement, they were the most abundant tree species in the region. However, conversion of natural areas to agriculture and development has removed many of the oaks from our region. Only 17% of oak ecosystems remain in the Chicago region. For more information on oak ecosystems in the Chicago region, see Chicago Wilderness's *Oak Ecosystem Recovery Plan*.

Prior to Euro-American settlement, oaks made up 60% of the regions canopy. Currently, in West Chicago, they account for 31.4% of the canopy. Adding additional oak trees would improve habitat for the birds, insects and wildlife that depend on them. This is particularly important in West Chicago, as it is situated on lands that were oak ecosystems prior to development.

Restoring oak ecosystems is a major focus of CRTI. It's efforts include improving oak management in natural areas, and encouraging their use in municipal plantings. Many municipalities avoid oaks because foresters believe that they do poorly as street trees. CRTI strives to dispel these biases, and to teach foresters how oaks can be used effectively in urban areas. The expanded use of oaks can help increase species diversity, and continue the legacy of oaks in our region.

Municipal plantings are not the only place where oaks can be used. West Chiago might also consider working with residents and commercial land owners to increase oaks abundance across all land use types. This could be done through programs like tree give-aways, or handouts that describe the benefits of planting oaks. CRTI is creating documents that could help in this endeavor.

West Chicago Urban Forest Summary

Woody invasive species like European buckthorn and bush honeysuckles make up almost one in three trees in the region. These shrubs were introduced as ornamental specimens, but they have escaped cultivation. Birds eat the berries produced by buckthorn and honeysuckle, allowing the seeds to be dispersed into natural areas. Both genera are extremely disruptive to native plants and animals. They create dense thickets, and prevent other species from growing around them (Fig. 15). In natural areas, they are one of the leading contributors to reduced oak regeneration.

Woody invasives are the most abundant in Cook, DuPage and Lake Counties, but they are becoming problematic region-wide (Fig. 16). While there are very few woody invasives in West Chicago's inventory, these trees likely exist on private property, or on unmanaged land. It is imperative to remove buckthorn from all land uses, as the seeds can easily travel to natural areas. It is difficult to dictate plantings on private property, but educating residents can encourage them to remove it of their own accord. This could include signage explaining invasive removal on public property, or expansion of programs like Conservation@Home.

Figure 15: A buckthorn thicket. Note that no other species are growing beneath the buckthorn.



Figure 16: Woody invasive abundance across all counties. They are the most abundant in Lake and McHenry Counties, but is a threat region-wide.