

Burlington Urban Forest Summary



The Chicago Region Trees Initiative (CRTI) goal is that, by 2050, the Chicago Region will support and host

a healthier urban forest, comprised of a diversity of tree species and ages, appropriately distributed across land use types in the region. The forest will provide the region improved environmental, economic, and social benefits. In order to achieve that goal CRTI works with a wide variety of people who work with and manage trees. This document is intended to help municipalities understand their urban forest, and identify strategies that they can use to make it better.

The Chicago Region Trees Initiative, USDA Forest Service, American Forests, and the University of Vermont mapped land cover across the seven-county Chicago Region. This project not only identifies tree canopy, but also other green infrastructure including vegetation under 10 feet tall, bare soil and water; and gray infrastructure including buildings, roads and rail and other paved surfaces like sidewalks and parking lots (Fig. 1). Here after, these seven layers will be referred to as *land cover types*.

The *urban forest* is comprised of all of the trees in an urban setting, regardless of who owns or manages them. It is made up of street trees, forested natural areas and even the trees in resident's back yards. These trees are all included in the urban forest, because they all provide benefits that municipalities depend on. They improve air and water quality, reduce flooding and the urban heat island effect, and reduce energy use by shading buildings. Trees provide habitat for wildlife and improve residents' quality of life by reducing crime rates, increasing property value and boosting social cohesion in neighborhoods.

The magnitude of benefits that trees provide correlates with the size, structure and location of their canopy. Understanding the extent of tree canopy is critical for urban planning. Canopy maps can be used to quantify the benefits that their trees provide, identify where new plantings would have the greatest impact and to develop priorities and strategies for expanding the canopy.




Fig. 1: Comparison of satellite image and land cover map. Seven types of gray and green infrastructure are in the land cover map.

Burlington Urban Forest Summary





Fig 2: Kane County's current land cover (A), including 16% canopy cover. An additional 76% of the county is suitable for planting (B). Burlington currently has 6% canopy cover (C), and 91% of the land cover could potentially be converted to canopy (D).

Overall, 16% of Kane County is covered by tree canopy (Fig. 2). There is a lot of room for growth across the county. We can identify spaces where trees could potentially be planted by adding together the vegetation, bare soil and other paved surface land cover types, as these land cover types could be converted to canopy with minimal effort. In all, these land cover types make up 76% of the county's area, meaning that canopy cover could potentially be raised to 92% if all of these surface were converted to trees. It is important to note, that while these surfaces could theoretically be covered with canopy, it is not necessarily preferable. Agricultural fields and baseball diamonds are included as "plantable space," but few would agree that these are ideal sites to expand the forest canopy.

These land cover data can also describe canopy at the municipal scale. Burlington currently has 6% canopy cover, and could potentially increase their canopy to 98% (Fig. 2).

Burlington Urban Forest Summary



Canopy cover is not distributed evenly across the region, nor within municipalities. To better understand how land cover patterns vary, we can compare them across land use types, like residential, commercial or park properties. In Burlington, the highest percentage of canopy is found in residential and vacant properties (Fig 3). Agricultural properties and natural areas have the lowest canopy cover. As one might expect, transit areas have the largest proportion of roads, and residential and commercial land use types have an abundance of buildings. See Table 1 at the end of this report for more details.

Fig 3: Variations in land cover across land use types.

Burlington Urban Forest Summary



By combining vegetation, bare soil and other paved surface categories we can identify which land use types have the most room for growth. In Burlington, the highest proportions of plantable space are found in agricultural properties and natural areas (Fig. 4).





Fig 4: Current canopy and possible planting space across land use types.


Burlington Forest Canopy Summary



While natural areas have a high proportion of plantable space, they make up a relatively small area in Burlington (Fig. 5). The majority of its land is agricultural.

Fig 5: The majority of land is agricultural land use.

Burlington Urban Forest Summary



The agricultural land has the most area that could possibly be converted to canopy (Fig. 6). Targeting these areas could have the greatest impact in expanding the canopy. Agricultural properties can be more challenging to plant trees on, but some canopy expansions can be made along roadways and drainage areas, or along farm buildings. Additionally, ordinances can mandate that trees be planted if these sites are converted from agricultural use.

Fig 6: Agricultural land has the greatest potential for increasing the canopy.

Table 1: Summary of land cover across land use types.

	Tree canopy		Vegetation		Bare soil		Water		Buildings		Roads and rail		Other paved	
	Acres	Percent	Acres	Percent	Acres	Percent	Acres	Percent	Acres	Percent	Acres	Percent	Acres	Percent
Agriculture	156.6	4.0%	3712.5	95.2%	0.5	0.0%	17.6	0.5%	3.3	0.1%	0.1	0.0%	7.4	0.2%
Cemetery	0.2	13.2%	1.0	86.2%	0.0	0.0%	0.0	0.0%	0.0	0.0%	0.0	0.4%	0.0	0.1%
Commercial	12.1	17.2%	48.1	68.5%	1.6	2.2%	3.1	4.4%	2.6	3.7%	0.1	0.1%	2.7	3.9%
Institutional	0.7	10.4%	4.0	62.1%	0.0	0.0%	0.0	0.0%	0.9	13.3%	0.0	0.2%	0.9	14.0%
Natural area	9.8	5.0%	184.0	94.5%	0.0	0.0%	0.2	0.1%	0.1	0.0%	0.0	0.0%	0.6	0.3%
Park	2.7	19.7%	10.3	74.4%	0.4	2.7%	0.0	0.0%	0.0	0.2%	0.0	0.0%	0.4	2.9%
Residential	54.9	33.3%	82.6	50.1%	0.1	0.1%	6.5	3.9%	11.9	7.2%	0.0	0.0%	8.7	5.3%
Transit	17.9	11.2%	77.7	48.5%	0.1	0.0%	0.1	0.1%	0.4	0.2%	56.2	35.1%	7.9	5.0%
Utility	0.5	9.2%	4.7	87.7%	0.0	0.0%	0.0	0.5%	0.0	0.3%	0.0	0.0%	0.1	2.4%
Vacant	39.8	65.1%	17.9	29.3%	0.0	0.0%	3.2	5.2%	0.1	0.2%	0.0	0.0%	0.1	0.2%
Total abundance	255.3	6.5%	4125.0	90.5%	2.6	0.1%	27.5	0.7%	19.1	0.4%	56.4	1.2%	28.8	0.6%

Burlington Urban Forest Summary



Urban trees are extremely valuable. Research has allowed us to quantify the values that trees provide, and these values go far beyond the aesthetics that are readily recognized. For example:

- Urban trees save energy by reducing surface temperatures and shading buildings.
- They store carbon dioxide and remove pollutants from the air.
- They intercept stormwater and help reduce flooding.
- Residents preferentially buy properties that have more trees, meaning that trees increase property values.

The i-Tree suite of tools was developed by the US Forest Service. They allow users to calculate tree benefits at a variety of scales, from an individual tree, to entire tree inventories, to landscape scale assessments of canopy and hydrology. For more information on i-Tree tools and methodology visit iTreetools.org.

Figure 7 shows the benefits that all of the trees (including trees public and private property) in Burlington offer. These values were calculated with i-Tree Landscape. Each year, Burlington's trees provide the municipality with \$24,600 worth of benefits. These trees also store a lot of carbon, which is valued at an additional \$1,423,000.




Figure 7: Trees offer myriad benefits, including intercepting stormwater, improving air quality and removing carbon from the atmosphere.

Burlington Urban Forest Summary



Cities tend to be hotter than rural areas because buildings and pavement absorb the sun's energy and release it as heat. This is known as the urban heat island effect. High urban temperatures increase the use of energy within buildings. It can also cause a variety of health issues to residents, and extreme heat can even cause death. Trees help lower urban temperatures by shading built surfaces and through evaporative cooling. Urban areas that have more tree canopy tend to have lower surface temperatures (Figure 8). Planting more trees in parking lots and around buildings can be especially helpful in reducing urban temperatures and making cities more comfortable.




Figure 8: The image on the left shows tree canopy and on the right shows surface temperature. Surface temperature was calculated using a landsat image from September 2014. Areas that have higher tree canopy tend to have lower temperatures.

Burlington Urban Forest Summary



Burlington has similar canopy to all of its neighbors except Campton Hills, which has considerably more (Fig. 9).





Figure 9: Comparison of land cover of Burlington and its neighbors.

**CHICAGO
REGION
TREES
INITIATIVE**

Our Trees.
Our Communities.
Our Future.



Municipal plantings are not the only place where oaks can be used. Municipalities might also consider working with residents and commercial land owners to increase oaks abundance across all land use types. This could be done through programs like tree give-aways, or handouts that describe the benefits of planting oaks. CRTI is creating documents that could help in this endeavor.

Burlington Urban Forest Summary



Woody invasive species like European buckthorn and bush honeysuckles make up almost one in three trees in the region. These shrubs were introduced as ornamental specimens, but they have escaped cultivation. Birds eat the berries produced by buckthorn and honeysuckle, allowing the seeds to be dispersed into natural areas. Both genera are extremely disruptive to native plants and animals. They create dense thickets, and prevent other species from growing around them (Fig. 11). In natural areas, they are one of the leading contributors to reduced oak regeneration.

Woody invasives are the most abundant in Cook, DuPage and Lake Counties, but they are becoming problematic region-wide (Fig. 12). While there are very few buckthorn and honeysuckle among municipality's managed trees, these trees likely exist on private property. It is imperative to remove buckthorn from all land uses, as the seeds can easily travel to natural areas. It is difficult to dictate plantings on private property, but educating residents can encourage them to remove it of their own accord. This could include signage explaining invasive removal on public property, or expansion of programs like Conservation@Home.



Figure 11: A buckthorn thicket. Note that no other species are growing beneath the buckthorn.




Figure 12: Woody invasive abundance across all counties. They are the most abundant in Lake and McHenry Counties, but is a threat region-wide.